

| Cripple | Creek | Catalog Mo | . 101-51-78 | -1/6 |
|--|---|----------------------|--|--|
| Catitude N. | 55° 24' 22' ' 4'' 31° 2' 24 '' 25'' | WR No. | 69 101-51- | 10 180 |
| Engeritande W. 1 | 31° 2' 24" 7 < " | n 1:0. | 09 | |
| Geodetic Map 1 | io. Ketchikan - B-4 | Work Area | Ketchikan | |
| Tocation N.W. o | corner Princess Bay | Watershed | Length 5 miles | |
| Drainage Area | of Watershed | | | |
| | Type Muskeg | | | |
| | | | | |
| Trails & Surve | y Routes Walkin | ng easy in lower st | ream - upper area look | s like |
| torrential fl | lows occur. | | | |
| Applal Survey | Notes Very limite | ed except at mouth | and I.T. area. | |
| | | | | |
| Anchorage Small | craft can enter lo | ower I.T. at low wa | ter. | |
| | | 1-16 of 61 and | | |
| Tide Stage Whe | en Surveyed <u>lst</u> | nail of 1100d. | | |
| FISHERY RESOUR | RCES | | | |
| Commercial I | Fisheries Pink sa | almon, a few coho c | bserverd 7-24-73. | |
| | | | | |
| Escapement | 1940, 1,000 pink, 2 | 200 chum; 1941, 10, | 000 pink,; 1947, 2 chu | im, 3 cono; |
| 1951, 3,700 p | oink, 55 chum, 1 red | 1; 1952, 80 pink,; | 1954, 2,000 pink, 95 c | hum; 1936, 200 |
| Spawning area: | 5435 m ² (3900'x30 |)'x50% ASA) | (pink, 150 c | .iium./ |
| Species Compos | ition: 90% pink, 1 | 0% chum. | | |
| Timing Late | . September, Octobe | er T lessen | | |
| genooting a | reas Limited in I | . I. lagoon. | | |
| Obs116ich D | of opriol Fair mu | iflats exposed at 1 | low tide. Shrimp larva | ae observed |
| in lower I.T | | dilats expeded at a | | |
| In tower 1.1 | riag Trout observe | ed, none taken in i | ry traps. Many good | looking pools |
| above Station | n #1 | cu, none cancar in . | 2, 22, 22, 23, 23, 23, 24, 24, 24, 24, 24, 24, 24, 24, 24, 24 | |
| | Present None | | | ************************************** |
| tand 085 at | 1169610 | | And the second s | alle and the second of the sec |
| Edstory of | Land Use None | | | |
| · · | | | | |
| Rehabilitat | ion Potential | Major spawning area | as free from debris. | |
| | | | | |
| Colls Unst | abilities from mid | stations #1 and #2 | upstream. | |
| a management of the second | | .1 | | |
| The second secon | | | | |
| THE RESOURCES | | | | |
| , | | A 2 2 2 4 | | _ |
| The second secon | | | sh) on banks, estimate | 9 |
| | Number of dropping | | | 1 |
| | Number seen on tid | | | 00 |
| _ | Wamber seen up cre | | | 0 |
| | Number of broods s | | | 0 |
| Mallarûs | Number seen on tid | | | 0 |
| | lamber seen up cre | | | 0 |
| | Number of broods | | | |
| | Darker of broods | | at mouth) | 3 |
| this Ragion- | Harler seen along | | | 0 |
| ** | Number of nests se | | | 0 |
| | Daller sean at mor | | | 500 ft. |
| 110.50 | Estimate length a | | • | 30 ft. |
| | Estimate depth out | | tt filmfer | 0% |
| | - MAI STABB PARSTA | ou whit persent o | 14.400 | |

PEAK ESCAPEMENT RECORD

Princess Bay, N.W. corner 101-51-78

| 101-51-78 | 3 | | | |
|-----------|--------|------|---------------|------------------------------------|
| DATE | PINK | СНИМ | OTHER SPECIES | REMARKS |
| 9-14-40 | 1,000 | 200 | | Good. 4000 fish off mouth |
| 10-6-41 | 10,000 | | | |
| 10-12-46 | | | | Most salmon @ mouth of stream. |
| 10-4-47 | | 2 | 3 coho | |
| 9-15-51 | 3,200 | 55 | 1 red | 2-3000 off mouth. 500 dead pink. |
| 8-14-52 | 80 | | | |
| 9-19-54 | 1,600 | 75 | 1 | 400 dead pink, 20 dead chum. |
| 9-14-56 | 200 | 150 | | 200 pink at mouth. |
| 9-7-66 | 250 | | | Poor visibility. Stream hi & dark. |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | - 85 - | |
| | _ | | | |